

5 TITLE: Method for the production of tablets by pressing and  
tablets produced by the method.

TECHNICAL FIELD:

10 The present invention relates to a method for the  
production of tablets by pressing of tablet material which  
contains microorganisms.

PRIOR ART:

15 Tablets are usually produced by pressing of a pulverulent  
tablet mass in a suitable shape in a so-called tablet  
punching machine. The tablets may have different shape and  
be of different size and they may also be of different  
hardness dependent on the properties of the tablet mass and  
the pressure to which they are subjected during the  
20 punching of the tablets.

25 When the tablets are formed heat is developed as a result  
of the friction against the mould surfaces and the inner  
friction in the tablet mass. Since the tablets usually  
consist of chemicals and the temperature increase is not  
too high, this will not create any problem since the  
chemicals can resist this heat increase and also are cooled  
rapidly. However, some tablet masses contain living  
microorganisms, such as bacteria, which are sensitive to  
30 high temperatures and because of this some of these  
bacteria die during the tablet punching.

TECHNICAL PROBLEM:

35 Tablets which contain microorganisms, for instance in the  
form of bacteria, and which are intended to contain such  
organisms will lose a part of or all of their value when  
the microorganisms are destroyed during the tablet  
punching. This cannot be avoided by simply using a lower  
pressure on the conventional tablet mass and thereby  
40 creating a lower heat development since the tablet must be

subjected to a certain pressure so that it maintains its shape and is not crumbled. For known tablet masses it is not unusual that a reduction of the viability (survival) of the bacteria in the tablet is up to 80% and even more.

5

**SOLUTION:**

It has therefore always been a problem to be able to produce tablets which contain microorganisms in the form of bacteria with a lesser reduction of the viability from tablet mass to a complete tablet and therefore according to the invention a method has been obtained for the production of tablets by pressing of tablet material comprising living organisms, which is characterized in that the tablet material also contains oligosaccharides consisting of more than two monosaccharides.

10

15

According to the invention, it is suitable that the oligosaccharides consist of fructose oligosaccharides, preferably inulin.

20

According to the invention it is suitable that the oligosaccharides are present in an amount of 40-99.5 % by weight of the tablet material.

25

The tablet material according to the invention can suitably contain microorganisms consisting of lactic acid producing bacteria.

30

The invention also comprises tablets produced by the method according to the invention, which tablets contain oligosaccharides and microorganisms whereby the oligosaccharides suitably consist of fructose oligosaccharides, preferably inulin.

35

The tablets according to the invention may contain lactic acid producing bacteria as microorganisms and they may also

064737 29959450

contain other additives such as polysaccharides, for example microcrystalline cellulose and starch, as well as other additives such as calcium diphosphate.

5 DETAILED DESCRIPTION:

The tablets according to the invention comprise microorganisms, preferably lactic acid producing bacteria cultures known as probiotica, which are intended to normalise or balance bacterial flora being present in the stomach and the intestine of humans or animals, but they may also contain other types of bacteria.

By mixing oligosaccharides, preferably fructose oligosaccharides, in the tablet mass as a so-called supporting substance the tablet punching is facilitated, which makes it possible to punch tablets at a lower pressure and lower heat development at the same time as the hardness of the tablet is maintained. The brittleness of the tablet, the friability, is surprisingly not changed with the tablet mass according to the present invention.

Due to this new composition, the punching pressure for the tablet making maybe reduced by up to 50% compared to conventional tablet punching methods without any reduction of the friability. This friability according to the invention will be 0.3-0.5, which is to be compared with the reference values which are accepted according to GMP (Good Manufacturing Practice) which are within the range of 0.1-1.0. The friability is expressed in percent weight reduction of the tablets when they are rotated 100 revolutions in a standard testing machine.

The amount of oligosaccharides depends on different crystalline qualities but may suitably be 99.5-40 weight percent of the total tablet mass without admixing any other supporting substance. However, if desired, known supporting

substances such as calcium diphosphate, microcrystalline cellulose and starch may be added in a suitable small amount. A smaller addition of oligosaccharides can, however, give rise to a smaller difference with regard to the viability compared with tablet masses containing only conventional supporting substances.

The tablets according to the present invention have a lower hardness due to the lower punching pressure when the tablets are formed but an increased viability for the strain of bacteria, which makes every tablet more efficient than conventional tablets. By not pressing the tablets so hard the yield of tablets for a given amount of tablet mass will also increase.

The invention will be described more in detail below by means of two examples, of which Example 1 describes a method according to the present invention and Example 2 describes a method of conventional kind.

Example 1: recipe having an active substance and tablet filling material

Str. thermophilus & L. bulgaricus	50%
Bifidobacterium animalis	0.5%
L. plantaris	0.5%
Inulin (fructose oligosaccharides)	<u>49%</u>
	<u>100%</u>

Hardness: 2.75 kp

Friability: 0.3

Viability original granulate: 5E8 cfu/g

Viability tablet: 3E8 cfu/g

40% reduction of cfu (colony forming units)

100 - 40 = 60

Example 2: recipe having active substance and tablet filling material

100 - 40 = 60%

(2)

Str. thermophilus & L. bulgaricus	50%
Bifidobacterium animalis	0.5%
L. plantaris	0.5%

5	Calcium diphosphate	20%
	Microcrystalline cellulose	18%
	Starch	11%
		100%

10      Hardness: 5.5 kp      Friability: 0.3%

         Viability original granulate: 5E8 cfu/g

         Viability tablet: 1E8 cfu/g

         80% reduction of cfu (colony forming units)

As appears from the above examples, the friability is maintained unchanged with a value of 0.3 whereas the hardness has been decreased to 2.75 kp compared with 5.5 kp for the conventional method. The viability has increased from 1E8 cfu/g to 3E8 cfu/g according to the invention. The reduction of cfu from tablet mass to tablet during the tablet punching became only 40% according to the new method and 80% according to the conventional method.

Accordingly, the new method results in an increased  
25 maintained viability after tablet punching of up to 200%  
compared with conventional tablet fillers. The increased  
yield results in an appreciably better economy and quality  
improvement of the above products.

30      The invention is not limited to the embodiments shown above  
but can be varied in different ways within the scope of the  
claims.

## 5 CLAIMS:

1. Method for the production of tablets having high  
viability in the tablet by pressing tablet material  
containing living microorganisms,  
10 c h a r a c t e r i z e d i n that the tablet material  
also contains oligosaccharides.

2. Method according to claim 1  
c h a r a c t e r i z e d i n that the oligosaccharides  
15 are present in an amount of 40-99.5 percent by weight of  
the tablet material.

3. Method according to any of claims 1-2,  
c h a r a c t e r i z e d i n that the oligosaccharides  
20 consist of fructose oligosaccharides.

4. Method according to any of claims 1-3,  
c h a r a c t e r i z e d i n that the oligosaccharides  
25 consist of inulin.

5. Method according to any of claims 1-4,  
c h a r a c t e r i z e d i n that the microorganisms  
consist of lactic acid producing bacteria.

6. Tablets produced according to any of claims 1-5  
30 containing oligosaccharides and microorganisms.

7. Tablets according to claim 6,  
c h a r a c t e r i z e d i n that the oligosaccharides  
35 consist of fructose oligosaccharides.

8. Tablets according to any of claims 6-7,  
c h a r a c t e r i z e d i n that the oligosaccharides  
40 consist of inulin.

9. Tablets according to any of claims 6-8,  
c h a r a c t e r i z e d i n t h a t t h e m i c r o o r g a n i s m s  
consist of lactic acid producing bacteria.

5 10. Tablets according to any of claims 6-9,  
c h a r a c t e r i z e d i n t h a t t h e y a l s o c o n t a i n  
polysaccharides such as microcrystalline cellulose and  
starch as well as other additives such as calcium  
diphosphate.

10

0946567-12199